

PROOF OF THE KNS CONJECTURE : D_r CASE

CHUL-HEE LEE

ABSTRACT. We prove the Kuniba-Nakanishi-Suzuki (KNS) conjecture concerning the quantum dimension solution of Q -system of type D_r obtained by a certain specialization of classical characters of the Kirillov-Reshetikhin modules. We use the affine Weyl group symmetry of quantum dimensions. As a result, we explicitly obtain the positive solution of the level k restricted Q -system of type D_r which plays an important role in dilogarithm identities for conformal field theories.

1. INTRODUCTION

1.1. Notations. Let $\mathfrak{g}$ be a simple Lie algebra of rank r and $\mathfrak{h}$ a Cartan subalgebra. $\mathfrak{h}^*$ denotes the dual space of $\mathfrak{h}$ and we use the symbol $\langle \cdot, \cdot \rangle$ to denote the natural pairing between $\mathfrak{h}$ and $\mathfrak{h}^*$.

Let $\Phi \subset \mathfrak{h}^*$ be the root system with its Dynkin diagram X with the index set $I = \{1, 2, \dots, r\}$. We will assume that X is simply laced. $\mathcal{C}(X) = (a_{ij})$ denotes the Cartan matrix. Let Q be the root lattice and P be the weight lattice. We fix a set Φ^+ of positive roots with the set $\Pi = \{\alpha_i | i \in I\}$ of simple roots. We will write $\alpha > 0$ if $\alpha \in \Phi^+$. For $\alpha = \sum_{i=1}^r c_i \alpha_i \in \Phi$, we define its height as $\text{ht } \alpha = \sum_{i=1}^r c_i$.

Let $\theta = \sum_{i=1}^r a_i \alpha_i$ be the highest root where a_i denotes the Dynkin labels. We denote the Coxeter number by $h = 1 + \text{ht } \theta$. Let $(\cdot | \cdot)$ be the non-degenerate invariant symmetric bilinear form on $\mathfrak{h}$ and on $\mathfrak{h}^*$ normalize $(\cdot | \cdot)$ so that $(\theta | \theta) = 2$.


$Q^\vee$ denotes the coroot lattice which is $\mathbb{Z}$ -dual of the weight lattice P . We will choose the basis $\Pi^\vee = \{h_i \in \mathfrak{h} | i \in I\}$ of the coroot lattice so that $\langle \alpha_i, h_j \rangle = a_{ji}$. Let $\{\omega_i \in P | i \in I\}$ be the dual basis of P for $\Pi^\vee$ so that $\langle \omega_i, h_j \rangle = \delta_{ij}$ and we call the elements of this set fundamental weights. The weight $\rho = \sum_{i=1}^r \omega_i \in P$ is called the Weyl vector.

We have the group algebra $\mathbb{Z}[P]$ with $\mathbb{Z}$ -basis of elements of the form e^λ , $\lambda \in P$. The multiplication is given by

$$e^\lambda e^\mu = e^{\lambda + \mu}.$$

We can regard e^λ as a function defined on $\mathfrak{h}^*$ by $\mu \mapsto e^{2\pi i(\lambda | \mu)}$. For a dominant weight $\lambda \in P^+$, χ_λ denotes the character of an irreducible highest weight module with highest weight λ and will be regarded as a function on $\mathfrak{h}^*$ since it is given by a ratio of two elements in $\mathbb{Z}[P]$.

Date: October 2012, last modified on October 8, 2012.

FIGURE 1. Dynkin diagram of type D_r

1.2. Statement of the KNS conjecture.

Definition 1.1. Let X be a Dynkin diagram of type ADE . For a family of variables $\{Q_i^{(a)} | a \in I, i \in \mathbb{Z}^{\geq 0}\}$, consider recurrences given by

$$(1.1) \quad \left(Q_i^{(a)}\right)^2 = \prod_{b \in I} \left(Q_i^{(b)}\right)^{\mathcal{I}(X)_{ab}} + Q_{i-1}^{(a)} Q_{i+1}^{(a)}$$

where $\mathcal{I}(X)$ denotes the adjacency matrix of X . We call this system of recurrence equations the unrestricted Q -system of type X . Throughout the paper, we will use boundary conditions $Q_0^{(a)} = 1$ for all $a \in I$.

Let q be a non-zero complex number which is not a root of unity. The Kirillov-Reshetikhin (KR) modules form a special class of finite dimensional modules of the quantum affine algebra $U_q(\hat{\mathfrak{g}})$ and they are parametrized by $a \in I$, $i \in \mathbb{Z}^{\geq 0}$ and $u \in \mathbb{C}$. Since the quantized universal enveloping algebra $U_q(\mathfrak{g})$ is contained in $U_q(\hat{\mathfrak{g}})$ as a subalgebra, for a given KR module $W_i^{(a)}(u)$, we can get a finite dimensional $U_q(\mathfrak{g})$ -module $\text{res } W_i^{(a)}(u)$ by restriction.

An important fact relevant to us is that the classical characters $Q_i^{(a)}$ of $\text{res } W_i^{(a)}(u)$ for $a \in I$ and $i \in \mathbb{Z}^{\geq 0}$ satisfy the unrestricted Q -system [Nak03, Her06]. $Q_i^{(a)}$ can be expanded into a sum of characters of irreducible modules of $\mathfrak{g}$ as

$$(1.2) \quad Q_i^{(a)} = \sum_{\omega \in \Omega_i^{(a)}} \chi_\omega$$

where $\Omega_i^{(a)}$ is a certain set of dominant weights depending on a and i . For example, when $X = A_r$, we have $Q_i^{(a)} = \chi_{i\omega_a}$ for $a \in I$ and $i \in \mathbb{Z}^{\geq 0}$ and they satisfy the unrestricted Q -system of type A_r . For $X = D_r$, the expression is more complicated and it is given by

$$(1.3) \quad Q_i^{(a)} = \begin{cases} \sum \chi_{k_a \omega_a + k_{a-2} \omega_{a-2} + \dots + k_1 \omega_1} & 1 \leq a < r-1, a \equiv 1 \pmod{2}, \\ \sum \chi_{k_a \omega_a + k_{a-2} \omega_{a-2} + \dots + k_0 \omega_0} & 1 \leq a < r-1, a \equiv 0 \pmod{2}, \\ \chi_{m \omega_a} & a = r-1, r \end{cases}$$

where $\omega_0 = 0$ and the summation is over all nonnegative integers satisfying $k_a + k_{a-2} + \dots + k_1 = i$ for a odd and $k_a + k_{a-2} + \dots + k_0 = i$ for a even. For more information about such decompositions, see [KNS11, Section 13] and references therein.

Let $k \geq 2$ be an integer. We are interested in solutions of the Q -system satisfying the unit boundary conditions $Q_k^{(a)} = 1$ for all $a \in I$.

Definition 1.2. We call the resulting system of equations

$$(1.4) \quad \begin{cases} Q_0^{(a)} = 1 & a \in I \\ \left(Q_i^{(a)}\right)^2 = \prod_{b \in I} \left(Q_i^{(b)}\right)^{\mathcal{I}(X)_{ab}} + Q_{i-1}^{(a)} Q_{i+1}^{(a)} & 1 \leq i < k, a \in I \\ Q_k^{(a)} = 1 & a \in I \end{cases}$$

for variables $\left(Q_i^{(a)}\right)$ with $0 \leq i \leq k$ and $a \in I$ the level k restricted Q -system of type X .

It is known that there exists a special unique solution of it possessing positivity and some additional properties as follows :

Theorem 1.3. [Lee12, Theorem 5.3.6] *Let X be a Dynkin diagram of type ADE of rank r . There exists a unique solution $\mathbf{z} = (z_i^{(a)})$ of the level k restricted Q -system of type X satisfying $z_i^{(a)} > 0$ for $0 \leq i \leq k$ and $a \in I$. Moreover, it satisfies the following additional properties:*

- (1) (symmetry) $z_i^{(a)} = z_{k-i}^{(a)}$ for $0 \leq i \leq k$ and $a \in I$.
- (2) (unimodality) $z_{i-1}^{(a)} < z_i^{(a)}$ for $1 \leq i \leq m = \lfloor \frac{k}{2} \rfloor$ and $a \in I$ where $\lfloor x \rfloor$ is the floor function.

Definition 1.4. We call the solution $\mathbf{z} = (z_i^{(a)})$ with $0 \leq i \leq k$ and $a \in I$ characterized in Theorem 1.3 the positive solution of the level k restricted Q -system.

One motivation to study the level k restricted Q -system comes from Nahm's conjecture [Nah07] and it was studied in detail in [Nah07, Kee07, NK09]. Also its unique positive solution plays an important role in the context of dilogarithm identities for conformal field theories [Kir89, Nak11] which have been the subject of intensive study. Let us briefly explain this.

Definition 1.5. The Rogers dilogarithm function is defined by

$$L(x) = -\frac{1}{2} \int_0^x \frac{\log(1-y)}{y} + \frac{\log(y)}{1-y} dy$$

for $x \in (0, 1)$. We set $L(0) = 0$ and $L(1) = \pi^2/6$ so that L is continuous on $[0, 1]$.

Theorem 1.6. [Nak11] *Let X be a Dynkin diagram of type ADE of rank r and $\mathfrak{g}$ be a simple Lie algebra associated with X . Let $\mathbf{z} = (z_i^{(a)})$ be the positive solution of the level k restricted Q -system of type X . For $1 \leq i \leq k-1$ and $a \in I$, let*

$$x_i^{(a)} = \frac{\prod_{b \in I} (z_i^{(b)})^{\mathcal{I}(X)_{ab}}}{(z_i^{(a)})^2}.$$

The following dilogarithm identity holds :

$$(1.5) \quad \frac{6}{\pi^2} \sum_{a \in I} \sum_{i=1}^{k-1} L(x_i^{(a)}) = \frac{k \dim \mathfrak{g}}{h+k} - \text{rank } \mathfrak{g} = \frac{(k-1)hr}{h+k}$$

where h denotes the Coxeter number of $\mathfrak{g}$.

For a physical interpretation of the rational number on the right hand side of (1.5), see [KNS11, Theorem 5.2] and references therein.

Our goal is to construct the positive solution of the level k restricted Q -system of type X explicitly. To achieve it, one may use classical characters of the Kirillov-Reshetikhin modules. If we regard $Q_i^{(a)}$ as a sum of characters of a simple Lie algebra using the decomposition (1.2), we can specialize it at the element $\frac{\rho}{h+k} \in \mathfrak{h}^*$ so that for $a \in I$ and $i \in \mathbb{Z}^{\geq 0}$, we get

$$z_i^{(a)} = Q_i^{(a)}\left(\frac{\rho}{h+k}\right) = \sum_{\omega \in \Omega_i^{(a)}} \chi_{\omega}\left(\frac{\rho}{h+k}\right)$$

and they solve the unrestricted Q -system. We call it the quantum dimension solution of the Q -system as will be explained in Section 2. In [KNS11], it has been conjectured that this specialization gives the positive solution of the level k restricted Q -system and also satisfies some additional level truncation properties as follows :

Conjecture 1.7. [KNS11, Conjecture 14.2.] Let $z_i^{(a)} = Q_i^{(a)}\left(\frac{\rho}{h+k}\right)$ for $a \in I$ and $i \in \mathbb{Z}^{\geq 0}$. It satisfies the following properties :

- (1) (positivity) $z_i^{(a)} > 0$ for $0 \leq i \leq k$ and $a \in I$.
- (2) (symmetry) $z_i^{(a)} = z_{k-i}^{(a)}$ for $1 \leq i \leq k-1$ and $a \in I$.
- (3) (unit boundary condition) $z_k^{(a)} = 1$ for $a \in I$.
- (4) (unimodality) $z_{i-1}^{(a)} < z_i^{(a)}$ holds true for $1 \leq i \leq m = \lfloor \frac{k}{2} \rfloor$ and $a \in I$ where $\lfloor x \rfloor$ is the floor function.
- (5) (occurrence of 0) $z_{k+1}^{(a)} = z_{k+2}^{(a)} = \cdots = z_{k+h-1}^{(a)} = 0$ for $a \in I$.

We call Conjecture 1.7 the KNS conjecture. This is already known to be true for A_r case. In this paper, we will prove the following :

Theorem. The KNS conjecture is true for $X = D_r$. Moreover, $z_{k+h}^{(a)} = 1$ for $1 \leq a \leq r-2$ and $z_{k+h}^{(r-1)} = z_{k+h}^{(r)} = \pm 1$.

This result will be divided into several parts and will be proved in Theorem 4.2, Theorem 4.5 and Theorem 4.7. The most tricky part lies in proving the unit boundary condition. Once we prove it, many results follow from Theorem 1.3. The key ingredients of our proof are the affine Weyl group symmetry and extended Dynkin diagram symmetry of quantum dimensions of affine weights obtained by suitable affinizations of weights. Although these are well-known concepts, they have not been effectively employed to attack our problem.




FIGURE 2. Extended Dynkin diagram of type $D_r^{(1)}$ and its Dynkin labels.

We review necessary results about quantum dimensions in Section 2 and carry out preliminary calculations involving the affine Weyl group in Section 3. Then the main results are obtained in Section 4.

Acknowledgements. This work is an outgrowth of the author's doctoral research. I would like to thank my advisor Richard E. Borcherds. I wish to thank Tomoki Nakanishi for helpful discussions at the MSRI workshop on Cluster Algebras and Commutative Algebra. I gratefully acknowledge financial support from Samsung Scholarship.

2. REVIEW ON QUANTUM DIMENSION AND ITS PROPERTIES

A good reference for this section is [FMS97, Section 16.3]. A few notions are tailored to our purposes. Here we collect results stated in [Lee12, Chapter 6] without proofs.

Definition 2.1. Let $\hat{P}$ be the lattice generated by $\hat{\omega}_0, \hat{\omega}_1, \dots, \hat{\omega}_r$ and $\hat{P}_k = \{\sum_{i=0}^r \lambda_i \hat{\omega}_i \in \hat{P} \mid \sum_{i=0}^r a_i \lambda_i = k\}$ where $a_0 = 1$. For a weight $\omega = \sum_{i=1}^r \lambda_i \omega_i \in P$, we define its level k affinization as $\hat{\omega} = \sum_{i=0}^r \lambda_i \hat{\omega}_i$ so that $\hat{\omega} \in \hat{P}_k$.

Let $\alpha_0 = -\theta$ and $\hat{\alpha}_j = \sum_{i=0}^r (\alpha_j | \alpha_i) \hat{\omega}_i$. We define the fundamental reflections $s_0, s_1, \dots, s_r$ on $\hat{P}$ linearly by

$$s_i \hat{\omega}_j = \hat{\omega}_j - \delta_{ij} \hat{\alpha}_i$$

and they generate the affine Weyl group W . The signature of $w \in W$ will be denoted by $(-1)^{\ell(w)}$.

Definition 2.2. Let $\lambda \in P$ and $\hat{\lambda} \in \hat{P}_k$ be its level k affinization. The quantum dimension or q -dimension of $\hat{\lambda}$ is defined by

$$(2.1) \quad \mathcal{D}_{\hat{\lambda}} = \chi_{\lambda} \left(\frac{\rho}{h+k} \right) = \frac{\prod_{\alpha > 0} \sin \frac{\pi(\lambda + \rho | \alpha)}{h+k}}{\prod_{\alpha > 0} \sin \frac{\pi(\rho | \alpha)}{h+k}}.$$

Theorem 2.3. Let $\lambda = \sum_{i=1}^l \lambda_i \omega_i \in P^+$ be a dominant weight such that $\sum_{i=1}^l a_i \lambda_i \leq k$. For its level k affinization $\hat{\lambda} \in \hat{P}_k^+$, $\mathcal{D}_{\hat{\lambda}} > 0$.

Definition 2.4. Let $\hat{\rho} = \sum_{i=0}^r \hat{\omega}_i \in \hat{P}$. We define the shifted affine Weyl group action on the set $\hat{P}$ by

$$w \cdot \hat{\lambda} = w(\hat{\lambda} + \hat{\rho}) - \hat{\rho}$$

for each $w \in W$.

Theorem 2.5. *Let $\hat{\lambda} \in \hat{P}_k$ and $w \in W$. $\mathcal{D}_{w \cdot \hat{\lambda}} = (-1)^{\ell(w)} \mathcal{D}_{\hat{\lambda}}$. If $w \in W$ is an element of odd signature and $w \cdot \hat{\lambda} = \hat{\lambda}$, $\mathcal{D}_{\hat{\lambda}} = 0$.*

Now we look at the role of the symmetry of the extended Dynkin diagram.

Theorem 2.6. *Suppose that $\hat{\lambda}_1, \hat{\lambda}_2 \in \hat{P}_k$ are conjugate by an automorphism of the extended Dynkin diagram. Then $\mathcal{D}_{\hat{\lambda}_1} = \mathcal{D}_{\hat{\lambda}_2}$.*

Corollary 2.7. *Let $\lambda = k\omega_i \in P^+$ be a dominant weight where ω_i is a fundamental weight such that $\hat{\omega}_i$ is conjugate to $\hat{\omega}_0$ by an automorphism of the extended Dynkin diagram. For its level k affinization $\hat{\lambda} \in \hat{P}_k$, $\mathcal{D}_{\hat{\lambda}} = 1$.*

3. PRELIMINARY COMPUTATIONS

From now on, we will assume $X = D_r$. Now we can write

$$z_i^{(a)} = Q_i^{(a)}\left(\frac{\rho}{h+k}\right) = \sum_{\omega \in \Omega_i^{(a)}} \mathcal{D}_{\hat{\omega}}$$

where the summation is given by (1.3).

Proposition 3.1. *Let $a \in \{1, r-1, r\}$. The following properties hold :*

$$(3.1) \quad z_i^{(a)} > 0 \quad \text{for } 0 \leq i \leq k,$$

$$(3.2) \quad z_i^{(a)} = z_{k-i}^{(a)} \quad \text{for } 0 < i < k,$$

$$(3.3) \quad z_k^{(a)} = 1.$$

Proof. Note that $z_i^{(a)} = \mathcal{D}_{(k-i)\hat{\omega}_0 + i\hat{\omega}_a}$. (3.1) follows from Theorem 2.3. By Theorem 2.6, $\mathcal{D}_{(k-i)\hat{\omega}_0 + i\hat{\omega}_a} = \mathcal{D}_{i\hat{\omega}_0 + (k-i)\hat{\omega}_a}$ and thus we get (3.2). (3.3) is a consequence of Corollary 2.7. $\square$

Proposition 3.2. *The followings properties hold :*

$$(3.4) \quad z_{k+j}^{(1)} = 0 \quad \text{for } 1 \leq j \leq h-1,$$

$$(3.5) \quad z_{k+h}^{(1)} = 1.$$

Proof. To prove (3.4), we use the product formula (2.1) for the quantum dimension. First note that for each integer l such that $1 \leq l \leq h-1 = 2r-1$, there exists a positive root α such that $\text{ht } \alpha = l$ and $\alpha - \alpha_1 \geq 0$. When we use the product formula for the quantum dimension $\mathcal{D}_{(k-i)\hat{\omega}_0 + i\hat{\omega}_1}$, only those roots may contribute to the product in a non-trivial way. Moreover, the number of such roots is exactly $h = 2r-2$. For $i \geq 0$, we have

$$\mathcal{D}_{(k-i)\hat{\omega}_0 + i\hat{\omega}_1} = \prod_{\substack{\alpha > 0 \\ \alpha - \alpha_1 \geq 0}} \frac{\sin \frac{\text{ht } \alpha + i}{h+k}}{\sin \frac{\text{ht } \alpha}{h+k}}.$$

Since $\{\text{ht } \alpha + i \mid \alpha > 0, \alpha - \alpha_1 \geq 0\}$ is the same as the set of all the integers from $1+i$ to $(h-1)+i$, we can conclude that $z_i^{(1)} = \mathcal{D}_{(k-i)\hat{\omega}_0 + i\hat{\omega}_1} = 0$

if $k+1 \leq i \leq k+(h-1)$ since there always exists a root α such that $\text{ht } \alpha + i = h+k$.

Now we turn to (3.5). When $i = h+k$, we get

$$\mathcal{D}_{(k-i)\hat{\omega}_0+i\hat{\omega}_1} = \prod_{\substack{\alpha > 0 \\ \alpha - \alpha_1 \geq 0}} \frac{\sin \frac{\text{ht } \alpha + (h+k)}{h+k}}{\sin \frac{\text{ht } \alpha}{h+k}} = \prod_{\substack{\alpha > 0 \\ \alpha - \alpha_1 \geq 0}} \frac{-\sin \frac{\text{ht } \alpha}{h+k}}{\sin \frac{\text{ht } \alpha}{h+k}}.$$

Since this product is over an even number $h = 2r-2$ of terms, the final product equals to 1 and it proves $z_{k+h}^{(1)} = 1$. $\square$

In the rest of the section, we will prove that for $2 \leq a \leq r-2$, $z_m^{(a)} = z_{m+1}^{(a)}$ when k is odd and $z_{m-1}^{(a)} = z_{m+1}^{(a)}$ when k is even where $m = \lfloor \frac{k}{2} \rfloor$. If $2 \leq a \leq r-2$, we will denote the element $k_a\hat{\omega}_a + k_{a-2}\hat{\omega}_{a-2} + \cdots + k_2\hat{\omega}_2 + k_0\hat{\omega}_0 \in \hat{P}_k$ by $(k_a, k_{a-2}, \dots, k_2, k_0)$ when a is even and $k_a\hat{\omega}_a + k_{a-2}\hat{\omega}_{a-2} + \cdots + k_1\hat{\omega}_1 + k_0\hat{\omega}_0 \in \hat{P}_k$ by $(k_a, k_{a-2}, \dots, k_1, k_0)$ when a is odd. Note that

$$z_i^{(a)} = \sum_{\hat{\omega} \in \hat{\Omega}_i^{(a)}} \mathcal{D}_{\hat{\omega}}$$

where

$$(3.6) \quad \hat{\Omega}_i^{(a)} = \left\{ (k_a, k_{a-2}, \dots, k_2, k_0) \in \hat{P}_k \mid \begin{array}{l} k_a + k_{a-2} + \cdots + k_2 \leq i \\ k_a, k_{a-2}, \dots, k_2 \in \mathbb{Z}^{\geq 0} \end{array} \right\}$$

for even a such that $2 \leq a \leq r-2$ and

$$(3.7) \quad \hat{\Omega}_i^{(a)} = \left\{ (k_a, k_{a-2}, \dots, k_1, k_0) \in \hat{P}_k \mid \begin{array}{l} k_a + k_{a-2} + \cdots + k_1 = i \\ k_a, k_{a-2}, \dots, k_1 \in \mathbb{Z}^{\geq 0} \end{array} \right\}$$

for odd a such that $2 \leq a \leq r-2$.

Assume that the level k is odd. Then $m = \frac{k-1}{2}$.

Proposition 3.3. *If $2 \leq a \leq r-2$ and a is even, $z_m^{(a)} = z_{m+1}^{(a)}$.*

Proof. Note that $\hat{\Omega}_m^{(a)} \subseteq \hat{\Omega}_{m+1}^{(a)}$ and

$$\hat{\Omega}_{m+1}^{(a)} \setminus \hat{\Omega}_m^{(a)} = \left\{ (k_a, k_{a-2}, \dots, k_2, k_0) \in \hat{P}_k \mid \begin{array}{l} k_a + k_{a-2} + \cdots + k_2 = m+1 \\ k_a, k_{a-2}, \dots, k_2 \in \mathbb{Z}^{\geq 0} \end{array} \right\}.$$

Because

$$k_0 + 2(k_a + k_{a-2} \cdots + k_2) = k = 2m+1,$$

$\hat{\omega} = (k_a, k_{a-2}, \dots, k_2, k_0) \in \hat{\Omega}_{m+1}^{(a)} \setminus \hat{\Omega}_m^{(a)}$ implies $k_0 = -1$. So for any $\hat{\omega} \in \hat{\Omega}_{m+1}^{(a)} \setminus \hat{\Omega}_m^{(a)}$, $\mathcal{D}_{\hat{\omega}} = 0$ since $s_0 \cdot \hat{\omega} = \hat{\omega}$. Thus $z_{m+1}^{(a)} = z_m^{(a)}$. $\square$

Proposition 3.4. *If $2 \leq a \leq r-2$ and a is odd, $z_m^{(a)} = z_{m+1}^{(a)}$*

Proof. If $\hat{\omega} = (k_a, k_{a-2}, \dots, k_1, k_0) \in \hat{\Omega}_{m+1}^{(a)}$ has $k_1 = 0$, $\mathcal{D}_{\hat{\omega}} = 0$ since $k_0 = -1$ and then $s_0 \cdot \hat{\omega} = \hat{\omega}$.

Let

$$(\hat{\Omega}_{m+1}^{(a)})' = \left\{ (k_a, k_{a-2}, \dots, k_1, k_0) \in \hat{P}_k \mid \begin{array}{l} k_a + k_{a-2} + \dots + k_1 = m+1 \\ k_a, k_{a-2}, \dots, k_1 \in \mathbb{Z}^{\geq 0}, k_1 \geq 1 \end{array} \right\}$$

and

$$\hat{\Omega}_m^{(a)} = \left\{ (k_a, k_{a-2}, \dots, k_1, k_0) \in \hat{P}_k \mid \begin{array}{l} k_a + k_{a-2} + \dots + k_1 = m \\ k_a, k_{a-2}, \dots, k_1 \in \mathbb{Z}^{\geq 0} \end{array} \right\}.$$

Let us construct a bijection between $(\hat{\Omega}_{m+1}^{(a)})'$ and $\hat{\Omega}_m^{(a)}$. Define a map from $(\hat{\Omega}_{m+1}^{(a)})'$ to $\hat{\Omega}_m^{(a)}$ by

$$(k_a, k_{a-2}, \dots, k_1, k_0) \mapsto (k_a, k_{a-2}, \dots, k_0, k_1).$$

For $(k_a, k_{a-2}, \dots, k_1, k_0) \in (\hat{\Omega}_{m+1}^{(a)})'$,

$$k_0 + k_1 + 2(m+1 - k_1) = k_0 - k_1 + 2m + 2 = 2m + 1$$

implies $k_0 = k_1 - 1 \geq 0$. From $k_a + k_{a-2} + \dots + k_1 + k_0 = m$, we have $(k_a, k_{a-2}, \dots, k_0, k_1) \in \hat{\Omega}_m^{(a)}$.

Conversely, any element $(k_a, k_{a-2}, \dots, k_1, k_0) \in \hat{\Omega}_m^{(a)}$ satisfies

$$k_0 + k_1 + 2(m - k_1) = k_0 - k_1 + 2m = 2m + 1.$$

Thus $k_0 = k_1 + 1 \geq 1$. So the map is a bijection between $(\hat{\Omega}_{m+1}^{(a)})'$ and $\hat{\Omega}_m^{(a)}$.

Note $\mathcal{D}_{(k_a, k_{a-2}, \dots, k_1, k_0)} = \mathcal{D}_{(k_a, k_{a-2}, \dots, k_0, k_1)}$ by Theorem 2.6. Therefore, $z_m^{(a)} = z_{m+1}^{(a)}$. $\square$

Now assume that the level k is even. Let $m = \frac{k}{2}$.

Lemma 3.5. *Let a be even and $2 \leq a \leq r-2$. For $\hat{\omega} = (k_a, k_{a-2}, \dots, k_2, -2) \in \hat{P}_k$ such that $k_a + k_{a-2} + \dots + k_2 = m+1$ and $k_2 = 0$, $\mathcal{D}_{\hat{\omega}} = 0$.*

Proof. $(s_0 s_2 s_0) \cdot \hat{\omega} = \hat{\omega}$. $\square$

Lemma 3.6. *Let a be even and $2 \leq a \leq r-2$. For $\hat{\omega} = (k_a, k_{a-2}, \dots, k_2, -2) \in \hat{P}_k$ such that $k_a + k_{a-2} + \dots + k_2 = m+1$ and $k_2 \geq 1$, $\mathcal{D}_{\hat{\omega}} = -\mathcal{D}_{\hat{\omega}'}$ where $\hat{\omega}' = (k_a, k_{a-2}, \dots, k_2 - 1, 0)$.*

Proof. $s_0 \cdot \hat{\omega} = \hat{\omega}'$. $\square$

Proposition 3.7. *If $2 \leq a \leq r-2$ and a is even, $z_{m-1}^{(a)} = z_{m+1}^{(a)}$.*

Proof. Recall that

$$\hat{\Omega}_{m-1}^{(a)} = \left\{ (k_a, k_{a-2}, \dots, k_2, k_0) \in \hat{P}_k \mid \begin{array}{l} k_a + k_{a-2} + \dots + k_2 \leq m-1 \\ k_a, k_{a-2}, \dots, k_2 \in \mathbb{Z}^{\geq 0} \end{array} \right\}$$

and

$$\hat{\Omega}_{m+1}^{(a)} = \left\{ (k_a, k_{a-2}, \dots, k_2, k_0) \in \hat{P}_k \mid \begin{array}{l} k_a + k_{a-2} + \dots + k_2 \leq m+1 \\ k_a, k_{a-2}, \dots, k_2 \in \mathbb{Z}^{\geq 0} \end{array} \right\}.$$

Let us define disjoint subsets R, S, T of $\hat{\Omega}_{m+1}^{(a)}$ by

$$\begin{aligned} R &= \{(k_a, k_{a-2}, \dots, k_2, k_0) \in \hat{\Omega}_{m+1}^{(a)} \mid k_a + k_{a-2} + \dots + k_2 = m+1, k_2 = 0\}, \\ S &= \{(k_a, k_{a-2}, \dots, k_2, k_0) \in \hat{\Omega}_{m+1}^{(a)} \mid k_a + k_{a-2} + \dots + k_2 = m+1, k_2 \geq 1\}, \\ T &= \{(k_a, k_{a-2}, \dots, k_2, k_0) \in \hat{\Omega}_{m+1}^{(a)} \mid k_a + k_{a-2} + \dots + k_2 = m\}. \end{aligned}$$

For $\hat{\omega} \in R$, $\mathcal{D}_{\hat{\omega}} = 0$ by Lemma 3.5 and so $\sum_{\omega \in R} \mathcal{D}_{\hat{\omega}} = 0$. Now we want to prove $\sum_{\omega \in S \cup T} \mathcal{D}_{\hat{\omega}} = 0$. For $(k_a, k_{a-2}, \dots, k_2, k_0) \in \hat{P}_k$, we have $k_0 + 2(k_a + k_{a-2} + \dots + k_2) = k = 2m$. Thus if $(k_a, k_{a-2}, \dots, k_2, k_0) \in S$, $k_0 = -2$. If $(k_a, k_{a-2}, \dots, k_2, k_0) \in T$, $k_0 = 0$. We can define a bijection between S and T by

$$(k_a, k_{a-2}, \dots, k_2, -2) \mapsto (k_a, k_{a-2}, \dots, k_2 - 1, 0).$$

By Lemma 3.6, $\sum_{\omega \in S \cup T} \mathcal{D}_{\hat{\omega}} = 0$. Therefore,

$$z_{m+1}^{(a)} = \sum_{\omega \in \hat{\Omega}_{m+1}^{(a)}} \mathcal{D}_{\hat{\omega}} = \sum_{\omega \in \hat{\Omega}_{m+1}^{(a)} \setminus (R \cup S \cup T)} \mathcal{D}_{\hat{\omega}}.$$

From $\hat{\Omega}_{m-1}^{(a)} = \hat{\Omega}_{m+1}^{(a)}$ and $\hat{\Omega}_{m-1}^{(a)} = \hat{\Omega}_{m+1}^{(a)} \setminus (R \cup S \cup T)$, we get $z_{m+1}^{(a)} = z_{m-1}^{(a)}$. $\square$

Lemma 3.8. *Let a be odd and $2 \leq a \leq r-2$. For $\hat{\omega} = (k_a, k_{a-2}, \dots, 1, -1) \in \hat{P}_k$, $\mathcal{D}_{\hat{\omega}} = 0$.*

Proof. $s_0 \cdot \hat{\omega} = \hat{\omega}$. $\square$

Lemma 3.9. *Let a be odd and $2 \leq a \leq r-2$. For $\hat{\omega} = (k_a, k_{a-2}, \dots, 0, -2) \in \hat{P}_k$, $\mathcal{D}_{\hat{\omega}} = 0$.*

Proof. $(s_0 s_2 s_0) \cdot \hat{\omega} = \hat{\omega}$. $\square$

Proposition 3.10. *If $2 \leq a \leq r-2$ and a is odd, $z_{m-1}^{(a)} = z_{m+1}^{(a)}$*

Proof. For any $\hat{\omega} = (k_a, k_{a-2}, \dots, k_1, k_0) \in \hat{\Omega}_{m+1}^{(a)}$ with $k_1 = 0$ or $k_1 = 1$, $\mathcal{D}_{\hat{\omega}} = 0$ by Lemma 3.8 and Lemma 3.9.

Let

$$(\hat{\Omega}_{m+1}^{(a)})' = \left\{ (k_a, k_{a-2}, \dots, k_1, k_0) \in \hat{P}_k \mid \begin{array}{l} k_a + k_{a-2} + \dots + k_1 = m+1 \\ k_a, k_{a-2}, \dots, k_3 \in \mathbb{Z}^{\geq 0}, k_1 \geq 2 \end{array} \right\}.$$

So we can write $z_{m+1}^{(a)} = \sum_{\omega \in (\hat{\Omega}_{m+1}^{(a)})'} \mathcal{D}_{\hat{\omega}}$.

Let us define a map from $(\hat{\Omega}_{m+1}^{(a)})'$ to $\hat{\Omega}_{m-1}^{(a)}$ by

$$(k_a, k_{a-2}, \dots, k_1, k_0) \mapsto (k_a, k_{a-2}, \dots, k_0, k_1).$$

For $(k_a, k_{a-2}, \dots, k_1, k_0) \in (\hat{\Omega}_{m+1}^{(a)})'$,

$$k_0 + k_1 + 2(m+1-k_1) = k_0 - k_1 + 2m + 2 = 2m.$$

This implies $k_0 = k_1 - 2 \geq 0$ and we have $(k_a, k_{a-2}, \dots, k_0, k_1) \in \hat{\Omega}_{m-1}^{(a)}$.

Conversely, any element $(k_a, k_{a-2}, \dots, k_1, k_0) \in \hat{\Omega}_{m-1}^{(a)}$ satisfies

$$k_0 + k_1 + 2(m-1-k_1) = k_0 - k_1 + 2m - 2 = 2m.$$

Thus $k_0 = k_1 + 2 \geq 2$. So the given map is a bijection between $(\hat{\Omega}_{m+1}^{(a)})'$ and $\hat{\Omega}_{m-1}^{(a)}$.

Since $\mathcal{D}_{(k_a, k_{a-2}, \dots, k_1, k_0)} = \mathcal{D}_{(k_a, k_{a-2}, \dots, k_0, k_1)}$ by Theorem 2.6, $z_{m+1}^{(a)} = z_{m-1}^{(a)}$. $\square$

4. PROOF OF THE MAIN THEOREM

In this section, we prove the main theorem using the results obtained in the previous section.

Lemma 4.1. *Let $\mathbf{w} = (w_i^{(a)})$ be a solution of the level k restricted Q -system such that $w_i^{(a)} \neq 0$ for $0 \leq i \leq k$ and $a \in I$. If $w_1^{(a)} = Q_1^{(a)}$ for any $a \in I$ and $\{Q_i^{(a)} | a \in I, i \in \mathbb{Z}^{\geq 0}\}$ satisfies the unrestricted Q -system, then $w_i^{(a)} = Q_i^{(a)}$ for $0 \leq i \leq k$ and $a \in I$. In particular, $Q_k^{(a)} = 1$.*

Proof. This is a direct consequence of the recursion (1.1)

$$Q_{i+1}^{(a)} = \frac{(Q_i^{(a)})^2 - \prod_{b \in I} (Q_i^{(b)})^{\mathcal{I}(X)_{ab}}}{Q_{i-1}^{(a)}}.$$

$\square$

Theorem 4.2. $\mathbf{z} = (z_i^{(a)})$ satisfies the following properties :

- (1) (positivity) $z_i^{(a)} > 0$ for $0 \leq i \leq k$ and $a \in I$.
- (2) (symmetry) $z_i^{(a)} = z_{k-i}^{(a)}$ for $1 \leq i \leq k-1$ and $a \in I$.
- (3) (unit boundary condition) $z_k^{(a)} = 1$ for $a \in I$.
- (4) (unimodality) $z_{i-1}^{(a)} < z_i^{(a)}$ for $1 \leq i \leq m = \lfloor \frac{k}{2} \rfloor$ and $a \in I$.

Proof. To prove the unit boundary condition, we divide the cases into when k is odd and k is even.

Assume that k is odd and let $m = \frac{k-1}{2}$. For $0 \leq i \leq k$ and $a \in I$, let

$$w_i^{(a)} = \begin{cases} z_i^{(a)} & 0 \leq i \leq m \\ z_{k-i}^{(a)} & m < i \leq k \end{cases}.$$

Since we have $z_m^{(a)} = z_{m+1}^{(a)}$ for any $a \in I$ by (3.2), Proposition 3.3 and Proposition 3.4, $\mathbf{w} = (w_i^{(a)})$ must be a solution of the level k restricted Q -system. Since $w_1^{(a)} = z_1^{(a)}$ and $w_i^{(a)} > 0$, we can conclude that $z_i^{(a)} = w_i^{(a)}$ for $0 \leq i \leq m$ by Lemma 4.1. Especially, $z_k^{(a)} = 1$ for any $a \in I$.

Assume that k is even and let $m = \frac{k}{2}$. For $0 \leq i \leq k$ and $a \in I$, let

$$(4.1) \quad w_i^{(a)} = \begin{cases} z_i^{(a)} & 0 \leq i \leq m \\ z_{k-i}^{(a)} & m < i \leq k \end{cases}.$$

Since we have $z_{m-1}^{(a)} = z_{m+1}^{(a)}$ for $a \in I$ by (3.2), Proposition 3.7 and Proposition 3.10, $\mathbf{w} = (w_i^{(a)})$ must be a solution of the level k restricted Q -system. Since $w_1^{(a)} = z_1^{(a)}$ and $w_i^{(a)} > 0$, again by Lemma 4.1, we can conclude that $z_i^{(a)} = w_i^{(a)}$ for $0 \leq i \leq m$ and $a \in I$. This proves $z_k^{(a)} = 1$ for any $a \in I$.

Now we have established the unit boundary condition. So for any $k \geq 2$, $\mathbf{z} = (z_i^{(a)})$ is the positive solution of the level k restricted Q -system characterized in Theorem 1.3 and it follows that $z_{i-1}^{(a)} < z_i^{(a)}$ for $1 \leq i \leq m = \lfloor \frac{k}{2} \rfloor$ and $a \in I$. $\square$

Now we prove $z_{k+j}^{(a)} = 0$ for $1 \leq j \leq h-1$ and any $a \in I$.

Proposition 4.3. $z_{k+1}^{(a)} = 0$ for any $a \in I$.

Proof. Since $z_k^{(a)} = 1$ and $z_{k-1}^{(a)} \neq 0$ for all $a \in I$ by Theorem 4.2, the recursion (1.1) for $z_k^{(a)}$:

$$(z_k^{(a)})^2 = \prod_{b \in I} (z_k^{(b)})^{\mathcal{I}(X)_{ab}} + z_{k-1}^{(a)} z_{k+1}^{(a)}$$

implies $z_{k+1}^{(a)} = 0$. $\square$

Lemma 4.4. Let $\{Q_i^{(a)} | a \in I, i \in \mathbb{Z}^{\geq 0}\}$ be a solution of the unrestricted Q -system. The following condition

$$\begin{bmatrix} Q_{i-1}^{(a-1)} & Q_{i-1}^{(a)} & Q_{i-1}^{(a+1)} \\ Q_i^{(a-1)} & Q_i^{(a)} & Q_i^{(a+1)} \\ Q_{i+1}^{(a-1)} & Q_{i+1}^{(a)} & Q_{i+1}^{(a+1)} \end{bmatrix} = \begin{bmatrix} * & 0 & * \\ 0 & Q_i^{(a)} & * \\ * & * & * \end{bmatrix}$$

implies $Q_i^{(a)} = 0$ for $2 \leq a \leq r-2$ where $*$ denotes an arbitrary number. Similarly, the following condition

$$\begin{bmatrix} Q_{i-1}^{(r-2)} & Q_{i-1}^{(r-1)} & Q_{i-1}^{(r)} \\ Q_i^{(r-2)} & Q_i^{(r-1)} & Q_i^{(r)} \\ Q_{i+1}^{(r-2)} & Q_{i+1}^{(r-1)} & Q_{i+1}^{(r)} \end{bmatrix} = \begin{bmatrix} * & 0 & 0 \\ 0 & Q_i^{(r-1)} & Q_i^{(r)} \\ * & * & * \end{bmatrix}$$

implies $Q_i^{(r-1)} = Q_i^{(r)} = 0$.

Proof. Again, this is a consequence of the recursion (1.1) for $Q_i^{(a)}$:

$$(Q_i^{(a)})^2 = \prod_{b \in I} (Q_i^{(b)})^{\mathcal{I}(X)_{ab}} + Q_{i-1}^{(a)} Q_{i+1}^{(a)}.$$

In both cases, we get $(Q_i^{(a)})^2 = 0$. $\square$

Theorem 4.5. $z_{k+j}^{(a)} = 0$ for $1 \leq j \leq h-1$ and any $a \in I$.

Proof. Recall (3.4) that $z_{k+j}^{(1)} = 0$ for $1 \leq j \leq h-1$. Since $z_{k+1}^{(a)} = 0$ for any $a \in I$ by Proposition 4.3, we get $z_{k+2}^{(a)} = 0$ by applying Lemma 4.4. Repeated applications of Lemma 4.4 imply $z_{k+j}^{(a)} = 0$ for $1 \leq j \leq h-1$ and any $a \in I$ inductively. $\square$

For the following lemma, let us put $Q_i^{(0)} = 1$ for convenience.

Lemma 4.6. *Let $\{Q_i^{(a)} | a \in I, i \in \mathbb{Z}^{\geq 0}\}$ be a solution of the unrestricted Q -system. The following condition*

$$\begin{bmatrix} Q_{i-1}^{(a-2)} & Q_{i-1}^{(a-1)} & Q_{i-1}^{(a)} & Q_{i-1}^{(a+1)} \\ Q_i^{(a-2)} & Q_i^{(a-1)} & Q_i^{(a)} & Q_i^{(a+1)} \\ Q_{i+1}^{(a-2)} & Q_{i+1}^{(a-1)} & Q_{i+1}^{(a)} & Q_{i+1}^{(a+1)} \end{bmatrix} = \begin{bmatrix} * & 0 & 0 & * \\ 1 & 1 & Q_i^{(a)} & * \\ * & * & * & * \end{bmatrix}$$

implies $Q_i^{(a)} = 1$ for $2 \leq a \leq r-2$ where $*$ denotes an arbitrary number.

Proof. Look the recursion (1.1) for $Q_i^{(a-1)}$:

$$(Q_i^{(a-1)})^2 = Q_i^{(a-2)}Q_i^{(a)} + Q_{i-1}^{(a-1)}Q_{i+1}^{(a-1)}.$$

With the given assumption, we get $1^2 = 1 \cdot Q_i^{(a)} + 0$. This proves $Q_i^{(a)} = 1$. $\square$

Theorem 4.7. $z_{k+h}^{(a)} = 1$ for any $1 \leq a \leq r-2$ and $z_{k+h}^{(r-1)} = z_{k+h}^{(r)} = \pm 1$.

Proof. Since $z_{k+h}^{(1)} = 1$ by (3.5) and $z_{k+h-1}^{(a)} = 0$ for any $a \in I$ by Theorem 4.5, we get $z_{k+h}^{(a)} = 1$ for $1 \leq a \leq r-2$ by applying Lemma 4.6. The recursions (1.1) for $z_{k+h}^{(r-2)}$, $z_{k+h}^{(r-1)}$ and $z_{k+h}^{(r)}$ give the following system of equations :

$$\begin{cases} (z_{k+h}^{(r-2)})^2 = z_{k+h}^{(r-1)}z_{k+h}^{(r)} \\ (z_{k+h}^{(r-1)})^2 = z_{k+h}^{(r-2)}z_{k+h}^{(r)} \\ (z_{k+h}^{(r)})^2 = z_{k+h}^{(r-2)}z_{k+h}^{(r-1)} \end{cases}.$$

From $z_{k+h}^{(r-2)} = 1$, we get $z_{k+h}^{(r-1)} = z_{k+h}^{(r)} = \pm 1$. $\square$

Remark 4.8. Similar analysis as in the proof of Proposition 3.2 using the product formula (2.1) for the quantum dimension can be used to show $z_{k+h}^{(r-1)} = z_{k+h}^{(r)} = 1$ when $r \equiv 0, 1 \pmod{4}$ and $z_{k+h}^{(r-1)} = z_{k+h}^{(r)} = -1$ when $r \equiv 2, 3 \pmod{4}$.

Example 4.9. Let $X = D_5$ and $k = 4$. We express $z_i^{(a)}$ in terms of quantum dimensions for $a \in I$ and $1 \leq i \leq h + k = 12$. For $a = 1, 4, 5$, we get

$$\begin{bmatrix} z_1^{(1)} & z_1^{(4)} & z_1^{(5)} \\ z_2^{(1)} & z_2^{(4)} & z_2^{(5)} \\ z_3^{(1)} & z_3^{(4)} & z_3^{(5)} \\ z_4^{(1)} & z_4^{(4)} & z_4^{(5)} \\ z_5^{(1)} & z_5^{(4)} & z_5^{(5)} \\ z_6^{(1)} & z_6^{(4)} & z_6^{(5)} \\ z_7^{(1)} & z_7^{(4)} & z_7^{(5)} \\ z_8^{(1)} & z_8^{(4)} & z_8^{(5)} \\ z_9^{(1)} & z_9^{(4)} & z_9^{(5)} \\ z_{10}^{(1)} & z_{10}^{(4)} & z_{10}^{(5)} \\ z_{11}^{(1)} & z_{11}^{(4)} & z_{11}^{(5)} \\ z_{12}^{(1)} & z_{12}^{(4)} & z_{12}^{(5)} \end{bmatrix} = \begin{bmatrix} \mathcal{D}_{3\hat{\omega}_0+\hat{\omega}_1} & \mathcal{D}_{3\hat{\omega}_0+\hat{\omega}_4} & \mathcal{D}_{3\hat{\omega}_0+\hat{\omega}_5} \\ \mathcal{D}_{2\hat{\omega}_0+2\hat{\omega}_1} & \mathcal{D}_{2\hat{\omega}_0+2\hat{\omega}_4} & \mathcal{D}_{2\hat{\omega}_0+2\hat{\omega}_5} \\ \mathcal{D}_{\hat{\omega}_0+3\hat{\omega}_1} & \mathcal{D}_{\hat{\omega}_0+3\hat{\omega}_4} & \mathcal{D}_{\hat{\omega}_0+3\hat{\omega}_5} \\ \mathcal{D}_{4\hat{\omega}_1} & \mathcal{D}_{4\hat{\omega}_4} & \mathcal{D}_{4\hat{\omega}_5} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ \mathcal{D}_{4\hat{\omega}_1} & \mathcal{D}_{4\hat{\omega}_4} & \mathcal{D}_{4\hat{\omega}_5} \end{bmatrix}.$$

For $a = 2, 3$, we have

$$\begin{bmatrix} z_1^{(2)} & z_1^{(3)} \\ z_2^{(2)} & z_2^{(3)} \\ z_3^{(2)} & z_3^{(3)} \\ z_4^{(2)} & z_4^{(3)} \\ z_5^{(2)} & z_5^{(3)} \\ z_6^{(2)} & z_6^{(3)} \\ z_7^{(2)} & z_7^{(3)} \\ z_8^{(2)} & z_8^{(3)} \\ z_9^{(2)} & z_9^{(3)} \\ z_{10}^{(2)} & z_{10}^{(3)} \\ z_{11}^{(2)} & z_{11}^{(3)} \\ z_{12}^{(2)} & z_{12}^{(3)} \end{bmatrix} = \begin{bmatrix} \mathcal{D}_{4\hat{\omega}_0} + \mathcal{D}_{2\hat{\omega}_0+\hat{\omega}_2} & \mathcal{D}_{3\hat{\omega}_0+\hat{\omega}_1} + \mathcal{D}_{2\hat{\omega}_0+\hat{\omega}_3} \\ \mathcal{D}_{4\hat{\omega}_0} + \mathcal{D}_{2\hat{\omega}_2} + \mathcal{D}_{2\hat{\omega}_0+\hat{\omega}_2} & \mathcal{D}_{2\hat{\omega}_0+2\hat{\omega}_1} + \mathcal{D}_{2\hat{\omega}_3} + \mathcal{D}_{\hat{\omega}_0+\hat{\omega}_1+\hat{\omega}_3} \\ \mathcal{D}_{4\hat{\omega}_0} + \mathcal{D}_{2\hat{\omega}_0+\hat{\omega}_2} & \mathcal{D}_{\hat{\omega}_0+3\hat{\omega}_1} + \mathcal{D}_{2\hat{\omega}_1+\hat{\omega}_3} \\ \mathcal{D}_{4\hat{\omega}_0} & \mathcal{D}_{4\hat{\omega}_1} \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ \mathcal{D}_{4\hat{\omega}_0} & \mathcal{D}_{4\hat{\omega}_1} \end{bmatrix}.$$

REFERENCES

- [FMS97] Philippe Di Francesco, P. Mathieu, and D. Sénéchal, *Conformal field theory*, Graduate Texts in Contemporary Physics, Springer, 1997.
- [Her06] David Hernandez, *The Kirillov-Reshetikhin conjecture and solutions of T -systems*, J. Reine Angew. Math. **596** (2006), 63–87.
- [Kee07] Sinead Keegan, *Algebraic K -theory and partition functions in conformal field theory*, Ph.D. thesis, University College Dublin, 2007.
- [Kir89] A. N. Kirillov, *Identities for the rogers dilogarithm function connected with simple lie algebras*, Journal of Mathematical Sciences **47** (1989), 2450–2459.
- [KNS11] Atsuo Kuniba, Tomoki Nakanishi, and Junji Suzuki, *T -systems and y -systems in integrable systems*, Journal of Physics A: Mathematical and Theoretical **44** (2011), no. 10, 103001.

- [Lee12] Chul-hee Lee, *Algebraic structures in modular q -hypergeometric series*, Ph.D. thesis, University of California, Berkeley, 2012.
- [Nah07] Werner Nahm, *Conformal field theory and torsion elements of the Bloch group*, Frontiers in number theory, physics, and geometry. II, Springer, Berlin, 2007, pp. 67–132.
- [Nak03] Hiraku Nakajima, *t -analogs of q -characters of Kirillov-Reshetikhin modules of quantum affine algebras*, Represent. Theory **7** (2003), 259–274 (electronic).
- [Nak11] Tomoki Nakanishi, *Dilogarithm identities for conformal field theories and cluster algebras: simply laced case*, Nagoya Math. J. **202** (2011), 23–43.
- [NK09] Werner Nahm and Sinead Keegan, *Integrable deformations of CFTs and the discrete Hirota equations*, ArXiv e-prints (2009).

MAX-PLANCK-INSTITUT FÜR MATHEMATIK, BONN, GERMANY
E-mail address: `chlee@mpim-bonn.mpg.de`